

UR-1542
 ER/40685/921
 hep-ph/9808403
 August 1998

Gluon Radiation in Top Quark Production and Decay at an e^+e^- Collider¹

Cosmin Macesanu and Lynne H. Orr

*Department of Physics and Astronomy, University of Rochester
 Rochester, New York 14627-0171*

Abstract. We study the effects of gluon radiation on top production and decay processes at an e^+e^- collider. The matrix elements are computed without any approximations, using spinor techniques. We use a Monte Carlo event generator which takes into account the infrared singularity due to soft gluons and differences in kinematics associated with radiation in the production versus decay process. The calculation is illustrated for several strategies of top mass reconstruction.

INTRODUCTION

The study of the top quark is one of the most important goals for future high energy experiments. Top has several characteristics which set it apart from the other quarks, the most obvious being its large mass: at 175 GeV top is 35 times heavier than its partner the b . The large mass of the top quark has a number of interesting consequences. It makes $t\bar{t}$ loop corrections to electroweak processes important, so that a good measurement of m_t coupled with extremely precise values of the W mass will give limits on the Higgs mass. A heavy top may decay into supersymmetric (SUSY) particles, providing a way of testing predictions of SUSY models. Perhaps most important, large mass means that the top Yukawa coupling to the Higgs is large; actually it is close to unity. This means that top studies can offer insight into electroweak symmetry breaking, as well as into the fermion mass generation process.

Another consequence of the top quark's large mass is that its decay width is about 1.5 GeV, which means its lifetime is of order 10^{-24} seconds. In this short

¹⁾ Presented by C. Macesanu at 20th Annual MRST (Montreal-Rochester-Syracuse-Toronto) Meeting on High-Energy Physics, Montreal, 13-15 May 1998.

a time, the top quark doesn't have time to hadronize. As a consequence, the top can be completely described by perturbative QCD. Also, the spin information is transmitted to the decay products (b and a W^+). This allows us to get information about the top couplings from angular distributions of the resulting particles; differences between measured and SM values would indicate new physics.

The top was discovered in 1995 at the Fermilab Tevatron proton-antiproton collider. This machine is now being upgraded, and in 2000 the study of the top here will start again. Also, at the Large Hadron Collider precise measurements of the top parameters (mass, couplings, production cross-section) will be one of the priorities.

Besides studies at these machines, it would be interesting to study top at a high-energy electron-positron collider. The principal advantage of a lepton collider versus a hadronic collider is a much cleaner environment; the task of performing certain precision measurements will be much easier in the absence of large QCD backgrounds. In particular, studies at the $t\bar{t}$ threshold can be done at a lepton collider, but such studies are impossible at a hadron collider. Measurements of top quark couplings are especially difficult at a hadron collider. At an e^+e^- collider, however, besides smaller background, the possibility of using polarized electrons in the initial state will be an advantage; also, the coupling of the top to neutral currents (γ and Z) will be accessible. In general, the capabilities of a high energy lepton collider are complementary to those of a hadron collider.

To be able to perform these studies, though, a good theoretical understanding of top quark's production and decay processes is essential. An important issue is QCD corrections due to real or virtual gluon emission because of the large value of the strong coupling constant (≈ 0.1). In the following, we will be concerned with corrections due to the radiation of a real gluon.

CALCULATIONAL PROCEDURE

We are interested in top production and decay with emission of a gluon:

$$e^+e^- \rightarrow \gamma^*, Z^* \rightarrow t\bar{t}(g) \rightarrow bW^+\bar{b}W^-(g)$$

(the top decays into a $b - W$ pair with a branching ratio close to unity). The gluon radiation can take place during the $t\bar{t}$ production process or during the decay of either the t or $\bar{t}$. Studies of this subject have focused on a single portion of the entire process (such as corrections to production [2,3] or decay [4] only), or imposed approximations such as soft gluons [5] or intermediate on-shell top quarks and massless b quarks [6]. The results presented here are obtained using an exact computation of the entire matrix element for real gluon emission, including all spin correlations, top width, and b mass effects [1].

Let us briefly describe the procedure for obtaining these results. We start by evaluating the amplitude for the process; there are four diagrams which contribute, (see Fig. 1), so we will have:




FIGURE 1. Feynman diagrams for top quark production with gluon radiation

$$M = \frac{B}{P_{tg} * P_t} + \frac{BB}{P_t * P_{tg}} + \frac{T}{P_t * P_{tg} * P_t} + \frac{TB}{P_t * P_t * P_{tg}} \quad (1)$$

with

$$P_t = p_t^2 - m_t^2 + im_t \Gamma_t ; p_t = p_{W^+} + p_b \quad (2)$$

$$P_{tg} = p_{tg}^2 - m_t^2 + im_t \Gamma_t ; p_{tg} = p_{W^+} + p_b + p_g$$

and similar definitions for the denominators of $\bar{t}$ propagators.

The B , BB terms correspond to the gluon being radiated by the b , $\bar{b}$ quarks. The last two terms correspond to a gluon radiated by the t or $\bar{t}$, either in the production or decay stage. To separate these cases, we rewrite the propagator products as follows:

$$\frac{1}{P_t * P_{tg}} = \frac{1}{2p_g p_t} \left[\frac{1}{P_t} - \frac{1}{P_{tg}} \right], \quad \frac{1}{P_{\bar{t}} * P_{t\bar{g}}} = \frac{1}{2p_g p_{\bar{t}}} \left[\frac{1}{P_{\bar{t}}} - \frac{1}{P_{t\bar{g}}} \right]$$

and the amplitude can be written

$$M = A1 + A2 + A3 \quad (3)$$

with

$$A1 = \frac{1}{P_{tg} * P_t} \left[B - \frac{T}{2p_g p_t} \right], \quad A2 = \frac{1}{P_t * P_{tg}} \left[BB - \frac{TB}{2p_g p_{\bar{t}}} \right]$$

$$A3 = \frac{1}{P_t * P_{\bar{t}}} \left[\frac{T}{2p_g p_t} + \frac{TB}{2p_g p_{\bar{t}}} \right]$$

We identify these three terms as corresponding to gluon radiation in the t decay (A1), $\bar{t}$ decay (A2) or production stage (A3).

The three separate parts of the amplitude are calculated using helicity amplitudes and can be evaluated numerically. We can then compute each of the six resulting terms separately:

$$\sum_{\text{helicities}} |M|^2 = \sum_{\text{helicities}} \left\{ |A1|^2 + |A2|^2 + |A3|^2 + 2\text{Re}[A1A2^* + A1A3^* + A2A3^*] \right\}$$

To integrate this formula over the phase space we use a three-channel Monte Carlo with one channel for each of the diagonal terms $|A1|^2, |A2|^2, |A3|^2$ and a combination of channels for the interference terms $2\text{Re}[A1A2^* + A1A3^* + A2A3^*]$. The phase space region where the gluon energy E_g goes to 0 presents some problems, though, because the amplitude has a singularity there. Even with cuts on E_g , the rapid variation of the integrand can spoil the integration procedure. To eliminate this problem, we tailor the momentum generator to the production of a gluon in association with two massive particles ($\gamma^*, Z^* \rightarrow t\bar{t}g$ or $t \rightarrow bWg$).

NUMERICAL RESULTS

In this section, we present some preliminary results obtained using the procedure described above. As input, we use the following values: $m_t = 175$ GeV, $\Gamma_t = 1.5$ GeV, $m_b = 5$ GeV, $\alpha_s = 0.1$, and a center of mass energy $W = 600$ GeV. Also, unless otherwise specified, we impose a cut on gluon energy $E_g > 10$ GeV.

First, we note that the radiation of a gluon plays an important role in $t\bar{t}$ production. The lowest-order, tree-level cross section is $\sigma_0 = 0.43$ pb, while the cross section for the process with emission of a gluon with energy greater than 10 GeV is $\sigma_0 = 0.39$ pb.² We see that the two quantities have the same order of magnitude.

An interesting (and important) issue is top mass reconstruction. This is closely related to the problem of assigning the gluon to the correct process in which it was radiated. For example, if the gluon was radiated in the production stage, the mass of the top is given by

$$m_t^2 = p_t^2 = (p_b + p_{W+})^2 \quad (4)$$

On the other hand, if the gluon was radiated in t decay stage, the above formula will give us a low estimate of m_t ; the correct formula will be

$$m_t^2 = p_{tg}^2 = (p_b + p_{W+} + p_g)^2 \quad (5)$$

²⁾ Note that this number includes contributions from decay-stage radiation which are not corrections to the total top production cross section.




FIGURE 2. Distributions in $\tilde{m}_t$ and $\tilde{m}_{tg}$ as defined in the text.

In Fig. 2 we present the mass distributions obtained using (4) and (5) respectively. While there are clear peaks at the input values of the top mass, we see significant tails due to wrong gluon assignments. The existence of such tails can increase measurement uncertainties on the top mass or confound attempts to identify top events by mass reconstruction. It is apparent that, for good mass reconstruction, we need to find a method to assign the gluon to the correct process in which it was




FIGURE 3. Mass distributions using mass cuts (left) and gluon angle cuts (right).

radiated.

Using the variables $\tilde{m}_t = \sqrt{p_t^2}$, $\tilde{m}_{tg} = \sqrt{p_{tg}^2}$, $\tilde{m}_{\bar{t}} = \sqrt{p_{\bar{t}}^2}$ and $\tilde{m}_{\bar{t}g} = \sqrt{p_{\bar{t}g}^2}$ we define four types of events:

- type 1 : $172 \text{ GeV} < \tilde{m}_{tg}, \tilde{m}_{\bar{t}} < 178 \text{ GeV}$
- type 2 : $172 \text{ GeV} < \tilde{m}_t, \tilde{m}_{\bar{t}g} < 178 \text{ GeV}$
- type 3 : $172 \text{ GeV} < \tilde{m}_t, \tilde{m}_{\bar{t}} < 178 \text{ GeV}$
- type 4 : any other event

We identify type 1 events as corresponding to gluons radiated in t decay stage. In this case we expect $p_{\bar{t}}$, p_{tg} to be on shell; however, as we use a Breit-Wigner distribution, we accept deviations from the exact value $m_t = 175 \text{ GeV}$ by about $2\Gamma_t = 3 \text{ GeV}$. Furthermore, we identify type 2 events as corresponding to gluons radiated in $\bar{t}$ decay stage and type 3 as corresponding to gluons radiated in $t\bar{t}$ production stage. As the gluon energy is quite big ($> 10 \text{ GeV}$) compared with Γ_t , we expect this interpretation to work; meaning that there will be very few events which satisfy one of the conditions 1, 2, 3 but which are open to more than one interpretation. Finally, type 4 would correspond to events for which there is no compelling evidence for either the production- or decay-stage radiation case; it will actually be a mixing of both.

In the first plot Fig. 3 we present the top mass distribution using events of types 1, 2, and 3 defined above. We have generated 250,000 events for each case. The smooth line is a Breit-Wigner function fitted over the mass distribution. The fit reproduces the input parameters with remarkable accuracy; it actually gives us




FIGURE 4. gluon-b angle distribution

175.04 GeV for m_t and 1.46 GeV for Γ_t .

Another method for reconstructing the process makes use of the gluon- b and gluon- $\bar{b}$ angle distributions. It is known that a gluon radiated by a quark tends to go in the same direction as that quark.³ Then, the gluons close to the b quark probably came from the top decay process; similarly, the gluons close to the $\bar{b}$ are likely to have come from $\bar{t}$ decay. We might guess that the rest of the gluons may be assigned to the production process.

In Fig. 4 we present the distribution of the angle θ_{bg} between the gluon and the b quark. The vertical-hatched part corresponds to type 1 events; and they are close to the b direction. The horizontal-hatched part corresponds to type 2 events; as they will cluster near the $\bar{b}$ quark, and b and $\bar{b}$ are mostly in opposite hemispheres, these events tend to gather at large angles. The cross-hatched part corresponds to type 3 events, and they are distributed uniformly. Finally, the non-hatched part corresponds to type 4 events, and, as they are an admixture of the first three, it looks like the whole distribution on a smaller scale.

Using this figure we can make the following conventions:

- if $\theta_{bg} < 0.7 \text{ rad}$ assign gluon to t decay
- if $\theta_{\bar{b}g} < 0.7 \text{ rad}$ assign gluon to $\bar{t}$ decay
- if $\theta_{bg}, \theta_{\bar{b}g} > 1 \text{ rad}$ assign gluon to $t\bar{t}$ production.

With these definitions, we construct the top mass distribution presented in the second plot in Fig. 3. Again, a fit with a Breit-Wigner function gives us values very close to our input parameters: $m_t = 175.05 \text{ GeV}$, $\Gamma_t = 1.6 \text{ GeV}$.

Fig. 5 presents the gluon energy distribution. The soft gluon singularity is visible (we have used a $E_g > 1 \text{ GeV}$ cut). This raises an interesting problem: soft gluons are not visible in the detector; therefore, the process in which a soft gluon is radiated will give the same signal as the process in which there is no gluon. The study of the experimental cross section for this last type of process will have to take into account radiation of gluons with energies up to 5 GeV, maybe. This also requires including virtual corrections, as in [6].

At this point, it is interesting to see if the interference terms contribute. We can integrate separately the contribution of diagonal terms and the contribution of interference terms; it turns out that this last contribution amounts only to around 1% of the total distribution, as shown in Fig. 5.

Up to here, we have worked at the parton level and presumed exact information about the momenta of particles in the final state. In a real experiment, though, there are a lot of complications: the electrons in the initial state can lose energy through radiation before interacting, thus having a lower CM energy; we don't see neutrinos, don't know which jet corresponds to which quark (or gluon), and the energies measured are not exact. One can expect that the detector resolution will have one of the biggest effect on the top mass (and width) reconstruction.

In Fig. 6 we present the distribution for the top mass (using gluon angle cuts)

³⁾ Since we keep the b mass, there is strictly speaking no collinear singularity; however gluons tend to be radiated from the b at small angles nonetheless.




FIGURE 5. Gluon energy distribution; total cross-section and interference contribution, respectively.




FIGURE 6. Mass reconstruction distribution, taking into account detector resolution (the dashed line corresponds to exact energy values).

obtained after taking into account this effect. The spread in the measured energies is parametrized by gaussians with widths $\sigma = 0.4\sqrt{E}$ for quarks and gluon, and $\sigma = 0.15\sqrt{E}$ for the W 's. The m_t value obtained in this case is still very good (175.03 GeV) but the width of the distribution in this case reflects rather the detector resolution effects than the top width.

CONCLUSIONS

We have presented preliminary results of a calculation of real gluon radiation in top production and decay. [1] Mass reconstruction using various mass cuts and gluon-bottom quark angle cuts was performed. In an ideal situation, (no initial state radiation, perfect particle identification, exact energy and angle measurements) the results were found to be very good, *i.e.*, the distributions faithfully reproduced the input mass and width. In a further study, we will have to take into account the effects of energy smearing due to detector resolution, initial state radiation, undetected neutrinos and jet identification problems. Effects due to interference between gluon radiation in the production and decay stages do not appear to be experimentally visible, at least for hard gluons. Soft gluon radiation (the experimental signal in this case will mimic that of a process without any gluon) and virtual gluon corrections have to be studied in more detail.

We thank C.R. Schmidt and W.J. Stirling for helpful correspondence and discussions. This work was supported in part by the U.S. Department of Energy and the National Science Foundation.

REFERENCES

1. C. Macesanu and L.H. Orr, in preparation.
2. J. Jersak, E. Laerman, and P. Zerwas, *Phys. Rev.* **D25** 1218 (1982)
3. Yu. L. Dokshitzer, V. A. Khoze, and W. J. Stirling, *Nucl. Phys.* **B428** 3 (1994)
4. M. Jezabek and J. Kuhn, *Phys. Lett.* **B329** 317 (1994)
5. V. Khoze, W. Stirling and L. Orr, *Nucl. Phys.* **B378** 413 (1992)
6. C. R. Schmidt, *Phys. Rev.* **D54** 3250 (1996)